

Medication Guide

ECZEMOL® Tablets

Read the Medication Guide that comes with **ECZEMOL®** before you start taking it and each time you get a prescription. There may be new information. This information does not take the place of talking with your doctor about your medical condition or your treatment.

What is **ECZEMOL®**?

ECZEMOL® is a biochemical homeopathic medication taken by mouth to treat moderate to severe eczema and atopic dermatitis. It has been found to work well with a variety of combination therapies.

Who should not take **ECZEMOL®**?

- **Do not** take **ECZEMOL** if you are pregnant, plan to become pregnant or become pregnant during **ECZEMOL®** treatment.
- Although there are no known contraindications, consult with your doctor prior to taking **ECZEMOL®** if you are allergic to any ingredients in **ECZEMOL®**. See the end of this Medication Guide for a complete list of ingredients in **ECZEMOL®**.

What should I tell my doctor before taking **ECZEMOL®**?

Tell your doctor if you have a history of kidney disease. The doctor will adjust the dosage.

Tell your doctor if you are pregnant.

Tell your doctor about all of the medicines you take including prescription and non-prescription medicines, vitamins and herbal supplements.

How should I take **ECZEMOL®**?

You must take **ECZEMOL®** exactly as prescribed by your doctor. It is important to take orally at the beginning of the day on an empty stomach, take nothing but water for one hour afterwards to avoid interference with **ECZEMOL®** absorption.

The amount of **ECZEMOL®** you take has been specially chosen for you. It is based upon your body weight.

ECZEMOL® may be swallowed whole, chewed or dissolved in the mouth.

If you miss a dose, just skip that dose. Do **not** take two doses at the same time.

If you take too much ECZEMOL® or overdose, call your doctor or poison control center right away.

What are the possible side effects of ECZEMOL®?

ECZEMOL® contains low doses of active ingredients. Therefore, there are minimal known side effects.

Tell your doctor if you have a history of contact sensitivity to nickel (common in metal) or if there is a history of vesicular hand eczema (dyshidrosis, pomphylx). Nickel sensitivity may be confirmed by a positive nickel patch test by your doctor. If you have nickel sensitivity, your doctor will prescribe a very low dose to start and then slowly increase your dosage over a 5 week period.

If you develop a rash, discontinue ECZEMOL® use and consult with your doctor.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088. You may also report side effects to Plymouth Pharmaceuticals, the manufacturer of ECZEMOL® at 1-800-316-9636.

How should I store ECZEMOL®?

- Store ECZEMOL® at room temperature, 68 to 77 Fahrenheit. Protect from light.
- Keep ECZEMOL® and all medicines out of the reach of children.

General Information about ECZEMOL®

Medicines are sometimes prescribed for conditions that are not mentioned in Medication Guides. Do not use ECZEMOL® for a condition for which it was not prescribed. Do not give ECZEMOL® to other people, even if they have the same symptoms that you have.

This Medication Guide summarizes the most important information about ECZEMOL®. If you would like more information, talk with your doctor.

You can ask your doctor or pharmacist for information about ECZEMOL® that is written for health care professionals. You can also visit: www.plymouthpharmaceuticals.com.

What are the ingredients in ECZEMOL®?

Active Ingredients: Potassium Bromide (Kali Bromatum) 1X, Sulphur 1X and Nickel Sulphate (Niccolum Sulphuricum) 1X. These drug ingredients are listed in the Homeopathic Pharmacopoeia of the United States (HPUS).

Inactive Ingredients: Lactose, Fumaric Acid and Magnesium Stearate.

ECZEMOL® is a registered trademark of Plymouth Healthcare Products LLC.

P.O. Box 702418

Tulsa, OK 74170-2418 USA

1-800-316-9636

info@plymouthpharmaceuticals.com

PCP#MG-002-10082014